

9501

REVISTA IBÉRICA de

# Aracnología

IA  
Grupo  
Ibérico de  
Aracnología

EA  
Asociación  
Ibérica de  
Aracnología


IB  
Instituto  
Ibérico de  
Aracnología



Instituto Ibérico de Aracnología / S.E.A.

# 9

ZARAGOZA, 30-VI-2004



ARTÍCULO:

**Notes on the collection of Salticidae (Araneae) from the Museum of Natural History 'Enrico Gaffi' in Bergamo, Italy**

Dmitri Logunov

The Manchester Museum, The University of Manchester, Oxford Road, Manchester M13 9PL, UK.  
dmitri.v.logunov@man.ac.uk

**Revista Ibérica de Aracnología**

ISSN: 1576 - 9518.

Dep. Legal: Z-2656-2000.

Vol. 9, 30-VI-2004

Sección: Artículos y Notas.

Pp: 271-275.

Edita:

**Grupo Ibérico de Aracnología (GIA)**

Grupo de trabajo en Aracnología de la Sociedad Entomológica Aragonesa (SEA)  
Avda. Radio Juventud, 37  
50012 Zaragoza (ESPAÑA)  
Tef. 976 324415  
Fax. 976 535697  
C-elect.: amellic@telefonica.net  
Director: A. Melic

Información sobre suscripción, índices, resúmenes de artículos *on line*, normas de publicación, etc. en:

Índice, resúmenes, abstracts vols. publicados:

<http://entomologia.rediris.es/sea/publicaciones/ria/index.htm>

Página web GIA:

<http://entomologia.rediris.es/gia>

Página web SEA:

<http://entomologia.rediris.es/sea>

**NOTES ON THE COLLECTION OF SALTICIDAE (ARANEAE) FROM THE MUSEUM OF NATURAL HISTORY 'ENRICO GAFFI' IN BERGAMO, ITALY**

Dmitri Logunov

**Abstract:**

Thirty-five salticid species were identified in the spider collection the Museo Civico di Scienze Naturali "Enrico Gaffi" (Bergamo, Italy), with new faunistic records from Croatia (2), Italy (30), Malta (1), Kenya (1), the Philippines (1) and Spain (2). *Saitis sanctaeufemiae* Kolosváry, 1938 is synonymized with *Saitis barbipes* (Simon, 1868). The female is described for *Leptorchestes mutilloides* (Lucas, 1846). Three faunistic records are new for particular regions: *Talavera monticola* (Kulczyński, 1884) for Italy, *Ergane carinata* Berry, Beatty et Prószyński, 1996 for the Philippines, and *Harmochirus luculentus* Simon, 1885 for Kenya.

**Key words:** Araneae, Salticidae, synonymy, faunistics, Italy, Croatia, Malta, Spain, Kenya, the Philippines.

**Notas sobre la colección de Salticidae (Araneae) del Museo de Historia Natural 'Enrico Gaffi' de Bergamo, Italia**

**Resumen:**

Han sido identificadas 35 especies de la colección aracnológica del Museo Cívico de Ciencias Naturales 'Enrico Gaffi' (Bérgamo, Italia), con nuevos registros para Croacia (2), Italia (30), Malta (1), Kenia (1), Filipinas (1) y España (2). *Saitis sanctaeufemiae* Kolosváry, 1938 es sinonimizado con *Saitis barbipes* (Simon, 1868). Se describe la hembra de *Leptorchestes mutilloides* (Lucas, 1846). Tres registros faunísticos son nuevos para regiones concretas: *Talavera monticola* (Kulczyński, 1884) para Italia, *Ergane carinata* Berry, Beatty et Prószyński, 1996 para Filipinas y *Harmochirus luculentus* Simon, 1885 para Kenia.

**Palabras clave:** Araneae, Salticidae, sinonimia, faunística, Italia, Croacia, Malta, España, Kenia, Filipinas.

**Introduction**

A small collection of Salticidae consisting of 104 adult specimens was loaned to me for identification by Dr P. Pantini of the Museo Civico di Scienze Naturali "Enrico Gaffi" (Bergamo, Italy; MBCG). A total of 35 species were identified, of which a list is provided below. This list is based on material mostly collected from Lombardia (23 of all the reported species) and it complements the preliminary list of the spiders of this region published recently by P. Pantini (2000); in the latter list (113 species in total) no salticid species were mentioned. Most of the identified salticid species from Lombardia can be characterized as common South-European, Euro-Central Asian or South-Palaearctic ones. The records of *Talavera monticola*, *Sitticus longipes* and especially the apparent female of *Leptorchestes mutilloides*, which is first to be found and described, are interesting and discussed in more detail below. For the rest of species the up-to-date and detailed information about their general distribution can be found in Metzner (1999) and Logunov & Marusik (2000). For a detailed account of the bibliography and records of Salticidae in Italy see Hansen (1985) and some subsequent works (e.g. Pesarini, 1995; Hansen, 1991; Alicata & Cantarella, 2000).

**A list of species***Aelurillus v-insignitus* (Walckenaer, 1802)

MATERIAL. ITALY: 3♂♂ (MBCG), Lombardia, LC, Vendrogno, S. da x Cammagiore - Betuleto, 1200 m a.s.l., 9.VI-6.VII.1999, P. Pantini; 1♀ (MBCG), Lombardia, Bergamo, Bonate sopra, River Brembo, grassland with bushes (pitfall traps), 195 m a.s.l., 13.IV-11.V.2002, S. Ciocca; 1♂2♀♀ (MBCG), Lombardia, Bergamo, Zanica, River Serio, arid grassland (pitfall traps), 190 m a.s.l., 16.IV-4.XI.2002, O. Lodovici, P. Pantini.

*Ballus chalybeus* (Walckenaer, 1802)

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, Bergamo, Astino, 270 m a.s.l., 30.V.2002, P. Pantini; 1♀ (MBCG), Lombardia, Bergamo, Zanica, River Serio, arid grassland (pitfall traps), 190 m a.s.l., 14.V-16.VI.2002, O. Lodovici, P. Pantini.

*Cyrba algerina* (Lucas, 1846)

MATERIAL. MALTA: 1♀ (MBCG), Radato, 9.V.2002, A. Mologni.

*Ergane carinata* Berry, Beatty et Prószyński, 1996

MATERIAL. THE PHILIPPINES: 1♀ (MBCG), SE of Luzon, III.1982, Pasquaré.  
COMMENTS. This species was recently described from Caroline Islands (Berry *et al.*, 1996) and hence the new locality in the Philippines is its westernmost record.

*Euophrys frontalis* (Walckenaer, 1802)

MATERIAL. ITALY: 2♂♂1♀ (MBCG), Lombardia, LC, Esino Lario, Ortanella, burnt thermophilous wood, 1100 m a.s.l., 7.IX-12.X.1999, P. Pantini, T.C.; 1♀ (MBCG), same locality, 10.XI.1999-13.III.2000, P. Pantini; 1♂ (MBCG), Emilia Romagna, PC, Bobbio, street toward Penice, 1100 m a.s.l., 20.VI-31.VII.2001, P. Pantini; 1♂1♀ (MBCG), Lombardia, Bergamo, Bonate Sotto, River Brembo, arid grassland (pitfall traps), 175 m a.s.l., 19.VI-8.VII.2002, S. Ciocca.

*Evarcha arcuata* (Clerck, 1758)

MATERIAL. ITALY: 1♂ (MBCG), Trentino-Alto Adige, Trento, Viotte del Monte Bondone, 1500 m a.s.l., 10.VIII.1998, P. Pantini.

*Evarcha falcata* (Clerck, 1758)

MATERIAL. ITALY: 1♂1♀ (MBCG), Toscana, FI, Marradi, Badia Valle, 430 m a.s.l., 1.X.2002, A. Usvelli; 1♀ (MBCG), Friuli Venezia Giulia, Pordenone, Barcis, Stream Cellina, 450 m a.s.l., 20.V.1996, P. Pantini, Valle; 1♂ (MBCG), Friuli Venezia Giulia, Pordenone, Claut, Stream Cellina, 950 m a.s.l., 16.IX.1996, P. Pantini, M. Valle.

*Evarcha jucunda* (Lucas, 1846)

MATERIAL. ITALY: 1♂ (MBCG), Basilicata, Potenza, Viggianello, pitfall traps, VII-VIII.1993, P. Pantini, M. Valle. - CROATIA: 1♀ (MBCG), Isola di Cres, Valun, at sea level, VIII.2001, P. Pantini.

*Harmochirus luculentus* Simon, 1885

MATERIAL. KENYA: 1♂ (MBCG), Nairobi, III.1981, M. Guerra.  
COMMENTS. This is a relatively widespread Afrotropical species, which is recorded from Kenya for the first time (cf. Logunov, 2001).

*Heliophanus auratus* C. L. Koch, 1835

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, LC, Cortenova, Stream Pioverna near Cortenova, 450 m a.s.l., 9.VI.1999, R. Manara, P. Pantini; 1♂ (MBCG), Lombardia, Bergamo, Bonate Sotto, River Brembo, arid grassland, 175 m a.s.l., 23.VII-9.VIII.2002, S. Ciocca; 1♂ (MBCG), Lombardia, Bergamo, Gromo, source, 700 m a.s.l., 2.VI.1994, R. Parrella, M. Valle; 1♀ (MBCG), Friuli Venezia Giulia, Pordenone, Vito d'Asio, source in the street towards Anduins, 270 m a.s.l., 21.V.1996, P. Pantini, M. Valle.

*Heliophanus cupreus* (Walckenaer, 1802)

MATERIAL. ITALY: 2♀♀ (MBCG), Liguria, SV, Bormida,

creek near Mt. Melogno, 670 m a.s.l., 28.V.2001, coll.?. 1♀ (MBCG), Liguria, SV, Calizzano, Mt. Quazzo, 800 m a.s.l., 29.V.2001, coll.?. 1♀ (MBCG), Lombardia, Bergamo, Carona, street towards Calvi, 1550 m a.s.l., 13.VII.1993, E. Bertuetti, M. Valle; 1♂ (MBCG), Lombardia, Bergamo, Endine Gaiano, tributary of Lake Gaiano, 340 m a.s.l., 7.VII.1999, P. Pantini, M. Valle; 1♀ (MBCG), Emilia Romagna, PC, Bobbio, street towards Penice, 1400 m a.s.l., 20.VI.2001, E. Ferrario, O. Lodovici, P. Pantini; 1♂ (MBCG), Lombardia, Bergamo, Caprino B.sco, S. Antonio d'Adda (source), 480 m a.s.l., 30.III.1994, E. Bertuetti; 1♀ (MBCG), Lombardia, Bergamo, Colzate, c/o Baite Sedernello, 1200 m a.s.l., 2.VIII.2001, P. Pantini *et al.*; 1♀ (MBCG), Lombardia, Bergamo, Zanica, River Serio, arid grassland (pitfall traps), 190 m a.s.l., 14.V-16.VI.2002, O. Lodovici, P. Pantini; 1♂ (MBCG), Lombardia, Bergamo, Valtorta, Fornonuovo, 900 m a.s.l., 16.VI.1993, E. Bertuetti, M. Valle; 1♀ (MBCG), Friuli Venezia Giulia, Udine, Resia, Uccia, 670 m a.s.l., 24.V.1996, P. Pantini, M. Valle; 1♂ (MBCG), Friuli Venezia Giulia, Pordenone, Vito d'Asio, street towards Anduins, 260 m a.s.l., 17.VII.1996, P. Pantini, M. Valle. - CROATIA: 1♀ (MBCG), Isola di Cres, Valun, on sea level, VIII.2001, P. Pantini.

*Heliophanus dubius* C. L. Koch, 1835

MATERIAL. ITALY: 3♂♂ (MBCG), Piemonte, CN, Briga Alta, Stream Negrone, 1600-1630 m a.s.l., 29-30.V.2001, coll.?.

*Heliophanus kochi* Simon, 1868

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, Bergamo, Solto Collina, San Defendente, grassland, 670 m a.s.l., 18.III.1997, P. Pantini, M. Valle; 1♀ (MBCG), Sardegna, SS, Olbia, Murta Maria, 28.VI.1993, C. Gusmini; 1♂ (MBCG), Lombardia, Bergamo, Bonate sopra, River Brembo, grassland with bushes (pitfall traps), 195 m a.s.l., 13.04-11.05.2002, S. Ciocca; 1♀ (MBCG), Lombardia, Bergamo, Zanica, River Serio, arid grassland (pitfall traps), 190 m a.s.l., 14.V-16.VI.2002, O. Lodovici, P. Pantini; 1♂ (MBCG), Abruzzo, Cherkio, Le Coste, 900 m a.s.l., 17.VI.1994, Stomelli.

*Heliophanus lineiventris* Simon, 1868

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Averara, Alpe Cul vers. Sud (pitfall traps), 1990 m a.s.l., 13.VIII-26.IX.2002, O. Lodovici, P. Pantini; 1♂ (MBCG), Lombardia, Bergamo, River Serio, not cultivated land (pitfall traps), 120 m a.s.l., 10.08-13.09.2002, S. Ciocca.

*Icius hamatus* (C. L. Koch, 1846)

MATERIAL. ITALY: 2♀♀ (MBCG), Lombardia, Bergamo, Dalmine, in a wasp nest, 20.VI.2000, Bugada.

*Icius subinermis* (Simon, 1937)

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, LO, Somaglia, Gargatano f. Po (pitfall traps), 45 m a.s.l., 3.IX.1997, F. Carrara, P. Pantini.

*Leptorchestes mutilloides* (Lucas, 1846) (Figs 3-4)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Predore, Corni, 550 m a.s.l., 14.V.2001, A. Soderi.  
DIAGNOSIS. In the arrangement and shape of the spermathecae, the female of *L. mutilloides* is most similar to *L. peresi* (Simon, 1868) (cf. Figs 3-4 and Figs 74-78 in Wesolowska & Szeremeta, 2001), but can easily be separated by the position of the copulatory pores directed laterally (anteriorly in *L. peresi*), by the shape of the glandular ducts (simple in *L. mutilloides* and multi-




Fig. 1-6. Copulatory organs of *Sitticus longipes* Canestrini, 1873 (1-2) from France (La Garve), *Leptorchestes mutilloides* (Lucas, 1846) (3-4) from Italy (Lombardia) and *Saitis sanctaeufemiae*, Kolosváry 1938 (= *Saitis barbipes* (Simon, 1868)) (5-6) from Italy (Calabria). 1, 4. Spermathecae, dorsal view. 2, 3. Epigyne, ventral view. 5, 6. Male palp, ventral and retrolateral views. Scale lines 0.1 mm.

chambered in *L. peresi*) and by the presence of two epigynal pockets in front of the epigastric furrow (absent in *L. peresi*).

COMMENTS. This female was provisionally assigned to *L. mutilloides*, which has so far been known only from males (s. Wesolowska & Szeremeta, 2001). The collected + does not belong to one of the six described *Leptorchestes* species recently revised by Wesolowska & Szeremeta (2001). From the latter revision, it is clear that the males of *L. mutilloides* are closest to those of *L. peresi*, as is the female described herein under the name of *L. mutilloides* (see "Diagnosis" above). Thus, although this ♀ was not collected together with a ♂, I am convinced it should belong to *L. mutilloides*.

DISTRIBUTION. S. Europe and the W. Mediterranean (Wesolowska & Szeremeta, 2001); rather common in Italy (see Hansen, 1985).

DESCRIPTION. *Female*. Carapace 2.05 long, 1.16 wide, 0.63 high at PLE. Ocular field 1.20 long, 0.90 wide anteriorly, 1.13 wide posteriorly. Diameter of AME 0.29. Chelicerae 0.83 long. Clypeus not marked. Abdomen 3.20 long, 1.60 wide. Leg segments:

I 1.03+0.53+0.83+0.48+0.48,  
II 0.78+0.48+0.55+0.41+0.33,

III 0.80+0.40+0.60+0.60+0.40,  
IV 1.23+0.55+0.98+0.93+0.48.

Leg spination. Leg I: tibia ventrally 1-2, metatarsus ventrally 2-2 apically. Leg II: tibia and metatarsus ventrally 1-0. Leg III without spines. Leg IV: metatarsus ventrally 1-0. Coloration. Carapace dark lustrously-brown, without covering scales. Sternum gray-brown. Labium, maxillae and chelicerae brown. Abdomen gray-brown, slightly constricted, with a white transverse line in the constricted area. Book-lung covers and spinnerets brown. Leg I: coxa yellow, femur brown+yellow, patella and tibia yellow, with a brown prolateral line, metatarsus and tarsus yellow. Leg II entirely yellow, with a brown prolateral line on all segments. Leg III: coxa and femur brown, patella and tibia yellow, with a brown prolateral line, metatarsus and tarsus entirely yellow. Leg IV: coxa, patella and metatarsus yellow+brown, femur and tibia entirely brown, tarsus entirely yellow. Epigyne and spermathecae as in Figs 3-4.

*Macaroeris nidicolens* (Walckenaer, 1802)

MATERIAL. ITALY: 2♀♀ (MMUM), Lombardia, Bergamo, Dalmine, in nido di vespa, 20.VI.2000, Bugada; 1♂ (MMUM), Lombardia, Bergamo, Valgoglio, source in wood, 800 m a.s.l., 2.VI.1994, Parrella, M. Valle.

*Marpissa muscosa* (Clerck, 1758)

MATERIAL. ITALY: 1♂1♀ (MBCG), Trentino Alto Adige, BZ, Ora, Stream Rio Nero, 250 m a.s.l., 6.IX.1997, P. Pantini, M. Valle.

*Mendoza canestrinii* (Ninni in Canestrini et Pavesi, 1868)

MATERIAL. ITALY: 1♂ (MBCG), Trentino Alto Adige, BZ, Montagna, Castelfeder, 335 m a.s.l., 6.IX.1997, P. Pantini, M. Valle.

*Menemerus semilimbatus* (Hahn, 1827)

MATERIAL. SPAIN: 1♂ (MBCG), Isola di Tenerife, Playa de Torviscas, 18.VIII.2002, O. Lodovici. – CROATIA: 1♀ (MBCG), Isola Bisevo, 22.VIII.2001, E. Ferrario.

*Myrmarachne formicaria* (De Geer, 1778)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Bariano, River Serio, uncultivated land (pitfall traps), 120 m a.s.l., 7-24.VII.2002, S. Ciocca.

*Pseudeophrys erratica* (Walckenaer, 1826)

MATERIAL. ITALY: 1♂ (MBCG), Piemonte, CN, Briga Alta, Lolle Traverso, 1630 m a.s.l., 30.V.2001, coll.?

*Pseudeophrys obsoleta* (Simon, 1868)

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, Bergamo, Bariano, River Serio, pebbly shore (pitfall traps), 120 m a.s.l., 14.V-16.VII.2002, S. Ciocca.

*Saitis barbipes* (Simon, 1868) (Figs 5-6)

*Attus barbipes* Simon, 1868: 563, pl. 6, f. 9 (D♂♀; the type series is in the Museum national d'Histoire naturelle, Paris, France; not examined). The species was reliably re-described and illustrated by Metzner (1999: plate 22).

*Saitis sanctaeufemiae* Kolosváry, 1938a: 303, f. 2, 4-5 (D♂♀; the syntypes in HNHM; examined). **New Synonymy.**

*Saitis sanctaeufemiae*: Kolosváry, 1938b: 583, f. 3 (♂♀). For a complete reference list see Platnick (2003).

MATERIAL. ITALY: 2♂ (HNHM, SALT360; the syntypes of *Saitis sanctaeufemiae*), Calabria, Strande des Meerbusens von Santa Eufemia, leg & det: Gábor Kolosváry; 4♀♀ (MBCG), Liguria, SV, Bormida, creek near Mt. Melogno, 670 m a.s.l., 28.V.2001; 2♂♂2♀♀ (MBCG), Lombardia, LC, Esino Lario, Ortanella, burnt thermophilous wood, 1100 m, 13.III-18.V.2000, P. Pantini; 1♀ (MBCG), Lombardia, LC, Vendrogno, Mornico, Castagneto, 970 m a.s.l., 13.III-18.V.2000, P. Pantini; 1♂2♀♀ (MBCG), Emilia Romagna, PC, Bobbio, wood on the pass Penice, 1100 m a.s.l., 20.V-20.VI.2001, P. Pantini, T.C.; 1♂1♀ (MMUM), same locality, 20.III-26.IV.2002, P. Pantini; 1♂ (MBCG), Lombardia, PV, S. Margherita di Staffora, beech wood near hotel Colletta, 1380 m a.s.l., 31.VII.2001, P. Pantini *et al.*; 1♀ (MBCG), Lombardia, PV, S. Margherita di Staffora, beech wood near hotel Colletta (pitfall traps), 1380 m a.s.l., 20.III-26.IV.2002, P. Pantini; 1♂3♀♀ (MBCG), same locality, 31.VII-19.IX.2001, P. Pantini; 2♀♀ (MBCG), Liguria, SV, Sassello, faggeta c/o Rio del Nido (pitfall traps), 1000 m a.s.l., 18.VII-10.X.2001, coll.?.; 1♂ (MBCG), Basilicata, Potenza, Viggianello, pitfall traps, VII-VIII.1993, P. Pantini, M. Valle.

COMMENTS. The species *S. sanctaeufemiae* was described from Italy (Calabria) from both sexes. As it is obvious from the original description (see Kolosváry, 1938a), that the main diagnostic characters used by the author to separate his new species from *Saitis barbipes* were

doubtful differences in leg spination in both sexes. The structure of the copulatory organs, though illustrated (1938a: fig. 4), was not taken into serious consideration. I re-examined the ♂ syntypes of *S. sanctaeufemiae* (Figs. 5-6), compared them to numerous comparative specimens of *S. barbipes* both from Italy and from other Mediterranean regions and came to the definite conclusion that the former name has to be synonymized with the latter one.

*Salticus scenicus* (Clerck, 1757)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Bariano, River Serio, pebbly shore (pitfall traps), 120 m a.s.l., 14.V-16.VII.2002, S. Ciocca.

*Sitticus distinguendus* (Simon, 1868)

MATERIAL. ITALY: 2♂♂ (MBCG), Lombardia, Bergamo, Bariano, River Serio, pebbly shore (pitfall traps), 120 m a.s.l., 7.VII-10.VIII.2002, S. Ciocca.

*Sitticus floricola* (C. L. Koch, 1837)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, BERGAMO, Endine Gaiano, tributary of Lake Gaiano, 340 m a.s.l., 7.VII.1999, P. Pantini, M. Valle.

*Sitticus longipes* Canestrini, 1873 (Figs 1-2)

MATERIAL. ITALY: 1♂1♀ (MMUM), Lombardia, Bergamo, Valbondione, Val Cerviera, 1900 m a.s.l., 11.VII.2002, R. Calandrino. – FRANCE: 1♀ (MBCG), La Grave, Glaciers de la Meije, 2800 m a.s.l., 11.VIII.2000, E. Ferrario.

COMMENTS. A rare species restricted to the high elevations of the Alps and recorded only from Switzerland, N. Italy, SE France and Austria (Tyrol) (s. Prószyński, 1973, 1976: map 188; Hansen, 1985; Pesarini, 1995; Thaler, 1997); previously recorded in France from two close localities by Simon (1937).

*Sitticus pubescens* (Fabricius, 1775)

MATERIAL. SPAIN: 1♂ (MBCG), Espot, Parque Nacional de Agues Tortes, Estany de Sant Maurici, 2000 m a.s.l., 16.VIII.2000, E. Ferrario.

*Sitticus rupicola* (C. L. Koch, 1837)

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, Bergamo, Mezzoldo, Val Terzera, 1500 m a.s.l., 22.VIII.1998, P. Pantini.

*Sitticus saltator* (O. Pickard-Cambridge in Simon, 1868)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Zanica, River Serio, arid grassland (pitfall traps), 190 m a.s.l., 16.VI-7.VII.2002, O. Lodovici, P. Pantini.

*Talavera aequipes* (O. Pickard-Cambridge, 1871)

MATERIAL. ITALY: 1♀ (MBCG), Lombardia, Bergamo, Bonate Sotto, River Brembo, arid grassland (pitfall traps), 175 m a.s.l., 23.VII-9.VIII.2002, S. Ciocca; 1♀ (MBCG), Lombardia, Bergamo, Bonate Sopra, River Brema, arid grassland (pitfall traps), 195 m a.s.l., 19.VI-8.VII.2002, S. Ciocca.

*Talavera monticola* (Kulczyński, 1884)

MATERIAL. ITALY: 1♂ (MBCG), Lombardia, Bergamo, Averara, Alpe Cul vers. Sud (pitfall traps), 1990 m a.s.l., 13.VIII-26.IX.2002, O. Lodovici, P. Pantini.

COMMENTS. This is a first record for Italy; to date, *T. monticola* has been known only from the mountains of Central Europe (the Alps, the Tatras and the Carpathians), where it occurs in (sub)alpine zone (s. Logunov & Kronstedt, 2003).

*Talavera* sp.

MATERIAL. ITALY: 1♀ (MBCG), 1♀ (MMUM), Lombardia, Bergamo, Bonate Sotto, River Brembo, arid grassland, 175 m a.s.l., 19.VI-8.VII.2002, S. Ciocca.

COMMENTS. This appears to be a new species. However, *Talavera* species are difficult to diagnose using isolated females (s. Logunov & Kronstedt, 2003). The taxonomic status of this species will be addressed elsewhere.

## Acknowledgements

I wish to cordially thank Dr P. Pantini for giving an opportunity to study the salticid collection under his care and for allowing me to keep some voucher specimens in the Manchester Museum (abbreviated in the text as MMUM). I am also obliged to Miss Galina Azarkina (Novosibirsk, Russia) for confirming the identification of *A. v-insignitus*. Miss Federica Menon (Manchester, UK) is thanked for her kind translating the Italian labels into the English and Mr Tamas Szuts (the Hungarian Natural History Museum, Budapest, Hungary; HNHM) for allowing to re-examine the syntypes of *Saitis sanctaeufemiae* Kolosváry, 1938. Dr J. Dunlop (Berlin, Germany) is much obliged for linguistic help.

## References

- ALICATA, P. & T. CANTARELLA 2000. I Salticidi di Sicilia: stato della conoscenza e descrizione di due nuove specie (Araneae Salticidae). *Mem. Soc. entomol. ital.*, **78**(2): 485-498.
- BERRY, J. W., J. A. BEATTY & J. PRÓSZYŃSKI 1996. Salticidae of the Pacific Islands. I. Distribution of twelve genera, with descriptions of eighteen new species. *J. Arachnol.*, **24**(3): 214-253.
- HANSEN, H. 1985. Contributo alla conoscenza dei Salticidae italiani (Arachnida: Araneae). *Boll. Mus. civ. St. nat. Venezia*, **34**: 241-322.
- HANSEN, H. 1991 (for 1989). Ricerche faunistiche del Museo Civico di Storia Naturale di Venezia nell'isola di Pantelleria. XI – Arachnida; Scorpiones, Pseudoscorpiones, Araneae. *Boll. Mus. civ. St. nat. Venezia*, **40**: 7-19.
- KOLOSVÁRY VON, G. 1938a. Eine neue Spinnenart: *Saitis Sanctae-Eufemiae* n. sp. *Zool. Anz.*, **121**: 303-306.
- KOLOSVÁRY VON, G. 1938b. Über calabrische Spinnen. *Festschr. Strand*, **4**: 582-585.
- LOGUNOV, D.V. 2001 (for 2000). A redefinition of the genera *Biamor* Peckham & Peckham, 1885 and *Harmochirus* Simon, 1885, with the establishment of a new genus *Sibianor* gen. n. (Araneae: Salticidae). *Arthropoda Sel.*, **9**(4): 221-286.
- LOGUNOV, D. V. & T. KRONESTEDT 2003. A review of the genus *Talavera* Peckham & Peckham, 1909 (Araneae, Salticidae). *J. Nat. Hist.*, **37**(9): 1091-1154.
- LOGUNOV, D. V. & YU M. MARUSIK 2000. *Catalogue of the jumping spiders of northern Asia (Arachnida, Araneae, Salticidae)*. Moscow: KMK Scientific Press. 299 pp.
- METZNER, H. 1999. Die Spingsspinnen (Araneae, Salticidae) Griechenlands. *Andrias*, **14**: 1-279.
- PANTINI, P. 2000. I Ragni del Sebino bergamasco (Italia, Lombardia) (Araneae). *Mem. Soc. entomol. ital.*, **78**(2): 361-378.
- PESARINI, C. 1995. Arachnida, Araneae. In: A. Minelli, Ruffo S. & La Posta S. (eds), *Checklist delle specie della fauna italiana*, 23. Calderini ed.: Bologna. 42pp.
- PLATNICK, N. 2003. The World Spider Catalog, Version 3.0, (Salticidae pages last updated December 29, 2003), American Museum of Natural History. <http://research.amnh.org/entomology/spiders/catalog81-87/index.html>
- PRÓSZYŃSKI, J. 1973. Revision of the genus *Sitticus* Simon, 1901 (Aranei, Salticidae), III. *Sitticus penicillatus* (Simon, 1875) and related forms. *Ann. Zool., PAN*, **30**(4): 71-95.
- PRÓSZYŃSKI, J. 1976. *Studium systematyczno-zoogeograficzne nad rodziną Salticidae (Aranei) Regionów Palearktycznego i Nearktycznego*. Wyższa Szkoła Pedagogiczna Siedlcach, No.6. S.1-260.
- SIMON, E. 1868. Monographie des espèces européennes de la famille des attides (Attidae Sundevall. - Saltigradae Latreille). *Ann. Soc. ent. Fr.*, (4) **8**: 11-72, 529-726.
- SIMON, E. 1937. *Les arachnides de France. Tome VI. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae; 5e et dernière partie*. Paris. Vol.6. Pp. 979-1298.
- THALER, K. 1997. Beiträge zur Spinnenfauna von Nordtirol – 4. Dionycha (Anyphaenidae, Clubionidae, Heteropodidae, Liocranidae, Philodromidae, Salticidae, Thomisidae, Zoridae). *Veröff. Mus. Ferdinandeum (Innsbruck)*, **77**: 233-285.
- WESOŁOWSKA, W. & M. SZEREMETA 2001. A revision of the ant-like salticid genera *Enoplomischus* Giltay, 1931, *Kima* Peckham & Peckham, 1902 and *Leptorchestes* Thorell, 1870 (Araneae: Salticidae). *Insect Syst. Evol.*, **32**: 217-240.